



Pictures Confirm Therapeutic Success:

Using SANUM preparations in Clinical Practice against Alopecia areata

by Dr. med. M. Al-Haj

In the course of their lives, many people will suffer from relatively sudden and severe localized hair loss, either once or repeatedly, the medical designation of which is Alopecia areata. This loss, usually localized at the outset to circular patches, frequently affects the head hair, and individual regions can gradually expand to total baldness. Other bodily regions can also be affected by Alopecia areata, such as armpit or facial hair. People with this disease often suffer, in fact, in the fullest sense of the word, when they feel themselves and their unlovely bald spots exposed to the gaze of the people around them. Any effective and lasting therapy is thus always gladly accepted by the afflicted patients.

At this point, orthodox medicine is still uncertain as to the etiology of this disease, but a familial connection has often been noted in this context, with many outbreaks occurring among family members. The possibility of it being an autoimmune disease has also been discussed. Besides the largely reversible conditions of this disease, there are also irreversible forms, such as Alopecia totale or Alopecia maligna. When the shed hair is examined closely, it turns out that the root end is not the normal club shape, but rather terminates in a sharp comma form. In this disease, the hair follicles are generally preserved. In treating this disease, orthodox medicine makes use of topical corticoids. In a departure from this practice, the biological SANUM therapy has successfully addressed the Alopecia problem using Professor Enderlein's ideas.



Fig. 1



Fig. 2




Fig. 3

The recent case described here concerns a 19 year old boy, who, in addition to pollen allergy, was diagnosed with Alopecia areata on his head. Figure 1 illustrates this partial hair loss. The success of the therapy was clearly visible after a mere four weeks. The course of therapy leading up to this success can be seen in Figures 2 through 4. The patient came in for treatment twice a week.



Fig. 4

The therapeutic measures and prescriptions consisted of the following in this case:

- In all, 10 i.m. compound injections with 0.5 ml of autologous blood twice weekly
 - + 1 ampule of PEFRAKEHL
 - + 1 ampule of ALBICANSAN
 - + 1 ampule of FORMASAN.
- Soft laser radiation twice weekly on the affected area.
- Intestinal cleansing as per the accompanying table.

- SANKOMBI 5X and PEFRAKEHL 5X drops: alternating daily, once a day rub 5 drops one at a time on the affected area.
 - SANUKEHL Strep 6X drops: take 10 drops every other day.
- Including intestinal cleansing, the therapy lasted about 12 weeks in all, of which the above measures took up just the first 5 weeks.

First published in the German language in the SANUM-Post magazine (38/1997)

© Copyright 1997 by Semmelweis-Institut GmbH, 27318 Hoya (Weser), Germany

All Rights Reserved

Intestinal Cleansing for Allergies During the Allergy Diet *Preliminary Therapy for Infection Susceptibility*

0. SANKOMBI drops
10 drops orally once a day before a meal for 2-3 weeks.

Therapy using Fungal Preparations

1. MUCOKEHL 5X ampules or drops
For 2 days: inject 1 ampule once at the acupuncture point Stomach 25, bilaterally and preperitoneally or i.v., or take 10 drops orally twice daily before meals.
2. FORTAKEHL 5X tablets or drops
For 20 days: either 1 tablet once daily before bed or 10 drops orally once daily before a meal.
3. PEFRAKEHL 5X capsules or drops
For 2-3 months: either 1 capsule once daily before bed or 10 drops orally once daily before a meal.

Therapy using Bacterial Preparations (for three weeks)

The following two preparations are administered alternately such that they are taken on different days of the week.

1. UTILIN 6X
Once, 1 capsule per week on an empty stomach without liquid for 1 week, then twice, 1 capsule per week on an empty stomach without liquid for 2 weeks.
2. RECARCIN capsules
Once, 1 capsule per week on an empty stomach with some liquid for 1 week, then twice, 1 capsule per week on an empty stomach with some liquid for 2 weeks.

Therapeutic recommendation by Dr. K. Werthmann (Salzburg; tested and approved by Dr. B. Köhler.)